

Drug Monograph

[Drug Name](#) | [Mechanism of Action and Pharmacokinetics](#) | [Indications and Status](#) | [Adverse Effects](#) | [Dosing](#) | [Administration Guidelines](#) | [Special Precautions](#) | [Interactions](#) | [Recommended Clinical Monitoring](#) | [Supplementary Public Funding](#) | [References](#) | [Disclaimer](#)

A - Drug Name

bevacizumab

SYNONYM(S): anti-VEGF monoclonal antibody; anti-VEGF rhuMAb

COMMON TRADE NAME(S): Avastin® (Hoffmann-LaRoche)

[back to top](#)

B - Mechanism of Action and Pharmacokinetics

Bevacizumab is a recombinant humanized monoclonal antibody that inhibits the biologic activity of human vascular endothelial growth factor (VEGF) and prevents binding of VEGF to its receptors on the surface of endothelial cells. VEGF binding initiates angiogenesis (endothelial proliferation and the formation of new blood vessels).

Absorption	Oral: Not orally bioavailable	
Distribution	Pharmacokinetics of bevacizumab are linear at doses ranging from 1 to 10 mg/kg. The predicted time to reach steady state is 100 days. Male subjects had a higher volume of distribution (+ 22%) than females.	
	Cross blood brain barrier?	No information found (unlikely)
	PPB	No information found
Metabolism	Bevacizumab is metabolized and eliminated via the reticuloendothelial system.	
	Active metabolites	no
	Inactive metabolites	no

Elimination

Males and patients with low albumin or high alkaline phosphatase have higher clearance (+20-26%).

Urine no

Half-life 20 days

[back to top](#)

C - Indications and Status**Health Canada Approvals:**

- In combination with fluoropyrimidine-based chemotherapy for first-line treatment of patients with metastatic carcinoma of the colon or rectum.
- In combination with carboplatin and paclitaxel for treatment of patients with unresectable advanced, metastatic or recurrent non-squamous non-small cell lung cancer.
- In combination with carboplatin and gemcitabine for treatment of patients with first recurrence platinum-sensitive epithelial ovarian, fallopian tube, or primary peritoneal cancer who have not received prior VEGF-targeted therapy.
- In combination with paclitaxel, topotecan, or pegylated liposomal doxorubicin for the treatment of patients with recurrent, platinum-resistant epithelial ovarian, fallopian tube or primary peritoneal cancer who received no more than two prior chemotherapy regimens and no prior VEGF-targeted therapy.

Health Canada Conditional Approvals

(pending the result of studies to verify the drug's clinical benefit. Patients should be advised of the nature of the marketing authorization granted.)

- As single agent for the treatment of patients with glioblastoma (WHO Grade IV) after relapse or disease progression, following prior therapy.

Notes:

Marketing approval for glioblastoma multiforme is conditional and based on trials demonstrating improvements in objective response, but not improvements in survival, symptoms or quality of life.

Approval for platinum-sensitive recurrent epithelial ovarian, fallopian tube or primary peritoneal cancer is based on an improvement in progression-free survival in patients with first recurrence after 6 months of platinum-based chemotherapy; no overall survival benefit was demonstrated.

Approval for platinum-resistant recurrent epithelial ovarian, fallopian tube or primary peritoneal cancer is based on an improvement in progression-free survival in patients with disease progression within 6 months of the most recent platinum-based chemotherapy; no overall survival benefit was found.

Other Uses:

- In combination with carboplatin and paclitaxel for treatment of patients with high risk epithelial ovarian, fallopian tube, or primary peritoneal cancer.
- In combination with platinum and paclitaxel for the treatment of advanced or recurrent cervical cancer.
- In combination with paclitaxel and topotecan for the treatment of advanced or recurrent cervical cancer in patients who cannot tolerate platinum-based chemotherapy.
- Renal Cell Cancer
- Mesothelioma

[back to top](#)

D - Adverse Effects

Emetogenic Potential: Minimal

Extravasation Potential: None

Adverse effects noted below are from pooled analyses of combinations of bevacizumab with chemotherapy where \geq grade 3 events were \geq 2% higher than chemotherapy alone (or any grade \geq 10% from pooled data or single trial). Exact incidences are specified where available.

ORGAN SITE	SIDE EFFECT* (%)	ONSET**
Cardiovascular	Arrhythmia (1-10% severe)	E
	<u>Arterial thromboembolism (4%)</u>	E
	Cardiotoxicity (\leq 13%)	I E
	<u>Hypertension (44%)</u>	I E
	Pulmonary hypertension	E D L
	<u>Venous thromboembolism 3-17% (8% severe)</u>	E
Dermatological	Hand-foot syndrome (1-10% severe)	E
	Other (cellulitis, 1-10%)	E
	Rash (\geq 10%) (may be severe)	E
Gastrointestinal	Abdominal pain (1-10% severe)	E
	Anorexia (\geq 10%)	E

	Constipation (≥10%) (severe ileus 1-10%)	E
	Diarrhea (≥10%) (severe)	I E
	GI obstruction (1-10%)	E
	GI perforation (1-10%; severe, includes gallbladder)	E
	Mucositis (≥10%)	E
	Nausea, vomiting (≥10%) (severe)	I E
General	Delayed wound healing (4%) /dehiscence	E
	Fatigue (≥10%)	E
	<u>Fistula (GI - 2%: nasal, tracheo-esophageal, biliary, urogenital; rare)</u>	E D L
Hematological	<u>Hemorrhage (≤50%) (5% severe)</u>	E
	INR / prothrombin time increased (grade 3/4- ≥2%)	E
	<u>Myelosuppression ± infection, bleeding (≥10%) (includes lymphopenia; severe)</u>	E
	Thrombotic microangiopathy (microangiopathic hemolytic anemia; rare)	E
Hypersensitivity	Hypersensitivity (≤5%)	I E
Infection	Infection (1-10% severe)	E
	Necrotizing fasciitis (rare)	E
Musculoskeletal	Musculoskeletal pain (≥10%)	E
	Osteonecrosis of jaw and other (rare)	E
Nervous System	Dysgeusia (≥10%)	E
	Headache (≥10%)	E
	Insomnia (20%)	E D
	<u>Neuropathy (≥10%) (severe)</u>	E
	<u>PRES (rare)</u>	E
	Seizure (rare)	E
	Somnolence (1-10% severe)	E
	Syncope (1-10%)	E
Ophthalmic	Eye disorders (10%)	E
Renal	Creatinine increased (<1%) (usually transient; may be severe)	E
	<u>Proteinuria (38%) (severe 10%)</u>	E D
Reproductive and breast disorders	Ovarian failure (39%)	D

Respiratory	Cough, dyspnea (≥10%)	E D
	Dysphonia 1-10%	E
	Pneumonitis (rare)	E D

* "Incidence" may refer to an absolute value or the higher value from a reported range.
 "Rare" may refer to events with < 1% incidence, reported in post-marketing, phase 1 studies, isolated data or anecdotal reports.

Dose-limiting side effects are underlined.

** I = *immediate* (onset in hours to days) E = *early* (days to weeks)
 D = *delayed* (weeks to months) L = *late* (months to years)

The most common side effects for bevacizumab include hemorrhage, hypertension, ovarian failure, proteinuria, insomnia, cardiotoxicity, anorexia, constipation, cough, dyspnea and diarrhea.

The most serious adverse effects include **gastrointestinal perforation, fistulas, hemorrhage, thromboembolism, severe hypertension (including PRES), cardiac and renal effects.**

Bevacizumab may exacerbate common toxicities of chemotherapy (hand foot syndrome, neurotoxicity, thrombocytopenia) when given in combination.

Hypertension is commonly observed, likely dose-dependent and should be managed with antihypertensives. Avoid diuretics in patients receiving cisplatin chemotherapy. The risk may be greater in platinum-sensitive, recurrent ovarian cancer patients. (For suggested treatment algorithm, see "Appendix 8: Management of Angiogenesis Inhibitor (AI) Induced Hypertension"). Pre-existing hypertension should be adequately controlled before starting bevacizumab treatment.

Patients with a history of hypertension may be at increased risk for the development of **proteinuria**. This is likely dose-related and may not completely resolve after discontinuing bevacizumab. Monitoring of proteinuria by dipstick urinalysis is recommended prior to starting and during bevacizumab therapy (see Monitoring section). Grade 4 proteinuria was seen in up to 1.4% of patients and was sometimes fatal.

Infusion and hypersensitivity reactions (including hypertension, respiratory and neurologic symptoms and hypertensive crisis) have been reported and appear to be more common when given in combination with chemotherapy.

Posterior reversible encephalopathy syndrome (PRES) is rare and may occur from 16 hours to 1 year after the start of bevacizumab. Patients present with seizures, headache, altered mental status, visual disturbance or cortical blindness, with or without associated hypertension. It is reversible if treated promptly after diagnosis with brain imaging, particularly MRI. Treatment of specific symptoms, including control of hypertension is recommended along with discontinuation of bevacizumab.

An increased risk of **arterial thromboembolic** events, including cerebrovascular accident, transient ischemic attack and myocardial infarction have been reported, especially in older patients, those with diabetes or prior events. **Venous thromboembolism** is also a risk, especially in patients with a prior history and patients with cervical cancer (increased risk reported in a clinical trial).

Congestive heart failure has been reported especially in metastatic breast cancer patients as well as patients with risk factors such as prior anthracyclines, radiation treatment, prior cardiovascular disease and in patients with lymphoma treated with R-CHOP. Bevacizumab is not authorized for the treatment of metastatic breast cancer.

Gastrointestinal perforation (including gallbladder) has been reported in all tumour types, although it is more common in colorectal and cervical cancer. An increased risk (nearly double) of intestinal perforation has been reported in colorectal cancer patients who have colorectal stents. The use of bevacizumab in these patients should be considered with caution (Health Canada review, Feb 2017).

Fistulas have been reported in all sites, including nasal septal perforation, and more common in colorectal cancer, especially with prior surgery or radiation in the area. Most fistulas were reported within the first 6 months of treatment, but may occur more than a year from treatment initiation. An increased risk (up to 8%) of gastrointestinal-vaginal fistulae was reported in patients with cervical cancer especially those patients who had received pelvic radiation. Bevacizumab is not approved for use in cervical cancer.

Hemorrhagic events may be life-threatening and include tumour-associated hemorrhage in all tumour types as well as CNS bleeding (especially in malignant glioma). The risk of hemoptysis is increased in patients with squamous NSCLC. There does not appear to be an increased risk in patients who are anticoagulated for venous thromboembolism.

Bevacizumab may adversely affect **wound healing** and should not be initiated for at least 28 days following major surgery (and held for 28 days prior to surgery, if elective) or until the surgical wound is fully healed.

Necrotizing fasciitis has been reported rarely, often secondary to wound healing complications, gastrointestinal perforation or fistulas. All patients were receiving additional chemotherapies other than bevacizumab; however, some patients did not have any other risk factors.

There is an increased risk of **myelosuppression** when used in combination with chemotherapy, and of thrombocytopenia when used in combination with gemcitabine/platinum-based treatment, especially in older patients.

Osteonecrosis of the jaw (ONJ) has been reported with prior or concomitant IV bisphosphonate treatment, radiation to the jaw, invasive dental procedures or glucocorticoid treatment.

Increased **ovarian failure** has been reported and appears to be reversible.

Intraocular inflammation or hemorrhage, retinal detachment or tear, increased need for cataract surgery, arterial thromboembolism and hypertension have been reported at increased rates with intraocular use, which is not approved in Canada and for which bevacizumab is not formulated.

Anti-bevacizumab and neutralizing antibodies have been observed rarely (<1%). The clinical significance of these is unknown.

[back to top](#)

E - Dosing

Refer to protocol by which patient is being treated. Bevacizumab should not be initiated in patients with recurrent hemoptysis, uncontrolled hypertension or wounds that require healing. Prior to treatment, a dental evaluation should be performed and major dental procedures completed. May consider hypersensitivity prophylaxis for patients who have had prior mild hypersensitivity reactions and are continuing on treatment.

Adults:

Continue bevacizumab until intolerable toxicity or disease progression.

Colorectal cancer:

Intravenous: 5 mg/kg Every 2 weeks

Glioblastoma:

Intravenous: 10 mg/kg Every 2 weeks

NSCLC:

Intravenous: 15 mg/kg Every 3 weeks

Recurrent epithelial ovarian, fallopian tube and primary peritoneal cancer (in combination with carboplatin and gemcitabine OR q3w topotecan):

Intravenous: 15 mg/kg Every 3 weeks

Recurrent epithelial ovarian, fallopian tube and primary peritoneal cancer (in combination with weekly topotecan, paclitaxel or pegylated liposomal doxorubicin in platinum-resistant patients):

Intravenous: 10 mg/kg Every 2 weeks

High Risk epithelial ovarian, fallopian tube and primary peritoneal cancer (in combination with carboplatin and paclitaxel):

Intravenous: 7.5 mg/kg Every 3 weeks

Dosage with Toxicity:

Dose reductions are not recommended. Bevacizumab should be held or discontinued based on toxicity.

Bevacizumab action	Toxicity		
	Any grade	Grade 3	Grade 4
Hold:	Uncontrolled hypertension		
	Delayed wound healing		
	Proteinuria \geq 2g/ 24 hours*		
	Surgery**		
Discontinue:		Hypertension (not controlled with medical management)	Hypertension
	Wound dehiscence, poor healing requiring medical intervention; necrotizing fasciitis		
	Nephrotic syndrome; non- recovery of proteinuria \geq 2g/24 hours		
	Severe hypersensitivity reaction		
	Tracheo-esophageal fistula, other non-GI fistulae		Any internal fistula
	GI perforation or fistula		
	PRES, hypertensive encephalopathy		
	Arterial thromboembolism	Pulmonary embolism	Venous thromboembolism (including pulmonary embolism)
	Symptomatic cardiac failure		

	Recurrent hemoptysis > 2.5 mL	Bleeding (any)	Bleeding (any)
	Intracranial bleeding		
<p>* may restart when < 2g/24hrs</p> <p>** for 28 days PRIOR (if surgery elective) and AFTER major surgery, or until wound healed</p>			

Dosage with Hepatic Impairment:

No information found. Not a major route of bevacizumab metabolism or excretion.

Dosage with Renal Impairment:

No information found. Not a major route of bevacizumab metabolism or excretion.

Dosage in the elderly:

Use with caution; patients > 65 years old have an increased risk of arterial thrombotic events as well as myelosuppression, fatigue, proteinuria, hypertension, dizziness, dysphonia, anorexia and GI effects.

Dosage based on gender:

Women have increased risk of severe hypertension, fatigue and abdominal pain, and lower bevacizumab clearance than males; however, no dose adjustment is required.

Children:

The safety and efficacy of bevacizumab in pediatric patients have not been studied. Osteonecrosis has been observed in clinical trials.

[back to top](#)

F - Administration Guidelines

- Bevacizumab infusions should not be administered or mixed with dextrose or glucose solutions due to potential for drug degradation.
- Mix in 100 mL bag NS. (Dilution should be 1.4 -16.5 mg/mL).
- Do not shake. Should not be mixed or diluted with other drugs.
- Compatible with PVC or polyolefin bags.
- DO NOT ADMINISTER AS AN IV PUSH OR BOLUS
- Infused over 90 minutes as loading dose, if tolerated next infusion can be given over 60 minutes; can thereafter be given over 30 minutes as maintenance dose
- Alternative infusion rates have been described by Mahfoud et al and Reidy et al, but these have not been approved by Health Canada
- Refrigerate unopened vials and protect from light; do not freeze.

[back to top](#)

G - Special Precautions

Contraindications:

- Patients with known hypersensitivity to bevacizumab or its components
- Chinese hamster ovary cell product or to other recombinant human or humanized antibodies
- Patients with untreated CNS metastases
- Patients with recurrent hemoptysis (>2.5ml), serious hemorrhage, or with squamous NSCLC
- Bevacizumab IV solution is not formulated for, and is contraindicated for intraocular use

Other Warnings/Precautions:

- Elderly patients
- Patients with a history of arterial thromboembolism or significant cardiovascular disease or cardiac failure
- Patients with coagulopathies (congenital, acquired or therapeutic)
- Hypertension should be controlled prior to starting treatment

- Bevacizumab should not be initiated for at least 28 days following major surgery or until wound healing has occurred; hold for 28 days prior to major elective surgery
- Use caution if given with bisphosphonates or other anti-angiogenic agents
- Do not use with diuretics in patients who are receiving platinum- based chemotherapy
- The safety and efficacy of concurrent radiotherapy and bevacizumab has not been established.
- Use with caution in colorectal cancer patients with colorectal stents; increased risk of GI perforation has been reported.

Other Drug Properties:

- Carcinogenicity: Unknown
Carcinogenicity and mutagenicity have not been studied.

Pregnancy and Lactation:

- Embryotoxicity: Yes
- Teratogenicity: Yes
Bevacizumab is not recommended for use in pregnancy. Cases of fetal abnormalities have been reported. Adequate contraception (including at least 2 contraceptive methods) should be used by both sexes during treatment, and for at least **6 months** after the last dose.
- Excretion into breast milk: Probable
Breastfeeding is not recommended.
- Fertility effects: Yes
Long-term effects unknown. Discuss fertility preservation with women of reproductive potential prior to starting treatment.

[back to top](#)

H - Interactions

No formal drug interaction studies have been performed. Combinations with EGFR monoclonal antibodies (e.g. cetuximab) and bevacizumab have not been studied and should not be administered for metastatic colorectal cancer. The safety and efficacy of concurrent radiotherapy has not been established.

AGENT	EFFECT	MECHANISM	MANAGEMENT
Anthracycline, thoracic radiation	↑ cardiotoxicity	Unknown	Caution
Irinotecan	Potential ↑ toxicity of Irinotecan	↑ SN38 (irinotecan metabolite) concentrations	Caution

sunitinib	Microangiopathic hemolytic anemia reported when used in combination	Unknown	Avoid this combination
Bisphosphonates, anti-angiogenic drugs	↑ risk of ONJ	Additive	Caution
Platinum or taxane-based chemotherapy	Increased risk of myelosuppression +/- infection, bleeding	Additive	Caution and closely monitor CBC

[back to top](#)

I - Recommended Clinical Monitoring

Treating physicians may decide to monitor more or less frequently for individual patients but should always consider recommendations from the product monograph.

Recommended Clinical Monitoring

Monitor Type	Monitor Frequency
CBC	Baseline and at each visit
Urine dipstick, 24 hour urine collection is recommended for patients with a 2+ or greater urine dipstick	Baseline and at each visit
Blood pressure	Baseline and every 2-3 weeks during therapy; more frequently in patients who develop hypertension
Dental evaluation	Baseline
Clinical assessment of hypersensitivity, perforation, fistula, GI symptoms, ONJ, hemorrhage, infection, myelosuppression, thromboembolism, delayed wound healing, hypertension, neurologic and cardiac effects	At each visit

Grade toxicity using the current [NCI-CTCAE \(Common Terminology Criteria for Adverse Events\) version](#)

Suggested Clinical Monitoring

Monitor Type	Monitor Frequency
Liver and renal function tests	Baseline and at each visit
Cardiac function tests (Echo, RNA and/or MUGA scans) especially in patients who are close to the lifetime cumulative dose of anthracyclines/anthracenediones	Baseline and as clinically indicated
INR for patients receiving warfarin	Baseline and regular

[back to top](#)

J - Supplementary Public Funding

New Drug Funding Program ([NDFP Website](#))

- Bevacizumab - First Line - Metastatic Colorectal Cancer
- Bevacizumab - Metastatic (Stage IVB), Persistent, or Recurrent Carcinoma of the Cervix
- Bevacizumab - Metastatic (Stage IVB), Persistent, or Recurrent Carcinoma of the Cervix
- Bevacizumab in combination with paclitaxel and carboplatin - Front-line Treatment Ovarian Fallopian Tube and Primary Peritoneal Cancer

[back to top](#)

K - References

Colorectal stents and bevacizumab: increased risk of intestinal perforation. Review. Health Canada Product InfoWatch. February 2017.

Dhillon, S. Bevacizumab combination therapy for the first-line treatment of advanced epithelial ovarian, fallopian tube or primary peritoneal cancer. *Drugs* 2012; 72(7): 917-30.

Gressett SM, Shah SR. Intricacies of bevacizumab-induced toxicities and their management. *Ann Pharmacother*. 2009 Mar;43(3):490-501.

Hurwitz H, Fehrenbacher L, Novotny W, et al. Bevacizumab plus irinotecan, fluorouracil, and leucovorin for metastatic colorectal cancer. *N Engl J Med* 2004;350:2335-42.

Mahfoud T, Tanz R, Mesmoudi M, et al. Bevacizumab 5 or 7.5 mg/kg in metastatic colorectal cancer can be infused safely over 10 minutes. *J Gastrointest Cancer*. 2012 Jun;43(2):244-8.

Product Monograph: Avastin® (bevacizumab). Roche Canada. February 2017.

Reidy DL, Chung KY, Timoney JP, et al. Bevacizumab 5 mg/kg can be infused safely over 10 minutes. *Journal of Clinical Oncology* 2007; 25: 2691-5.

March 2017 added warning re: increased risk of GI perforation in patients with colorectal stents

[back to top](#)

L - Disclaimer

Refer to the [New Drug Funding Program](#) or [Ontario Public Drug Programs](#) websites for the most up-to-date public funding information.

The information set out in the drug monographs, regimen monographs, appendices and symptom management information (for health professionals) contained in the Drug Formulary (the "Formulary") is intended for healthcare providers and is to be used for informational purposes only. The information is not intended to cover all possible uses, directions, precautions, drug interactions or adverse effects of a particular drug, nor should it be construed to indicate that use of a particular drug is safe, appropriate or effective for a given condition. The information in the Formulary is not intended to constitute or be a substitute for medical advice and should not be relied upon in any such regard. All uses of the Formulary are subject to clinical judgment and actual prescribing patterns may not follow the information provided in the Formulary.

The format and content of the drug monographs, regimen monographs, appendices and symptom management information contained in the Formulary will change as they are reviewed and revised on a periodic basis. The date of last revision will be visible on each page of the monograph and regimen. Since standards of usage are constantly evolving, it is advised that the Formulary not be used as the sole source of information. It is strongly recommended that original references or product monograph be consulted prior to using a chemotherapy regimen for the first time.

Some Formulary documents, such as the medication information sheets, regimen information sheets and symptom management information (for patients), are intended for patients. Patients should always consult with their healthcare provider if they have questions regarding any information set out in the Formulary documents.

While care has been taken in the preparation of the information contained in the Formulary, such information is provided on an "as-is" basis, without any representation, warranty, or condition, whether express, or implied, statutory or otherwise, as to the information's quality, accuracy, currency, completeness, or reliability.

CCO and the Formulary's content providers shall have no liability, whether direct, indirect, consequential, contingent, special, or incidental, related to or arising from the information in the Formulary or its use thereof, whether based on breach of contract or tort (including negligence), and even if advised of the possibility thereof. Anyone using the information in the Formulary does so at his or her own risk, and by using such information, agrees to indemnify CCO and its content providers from any and all liability, loss, damages, costs and expenses (including legal fees and expenses) arising from such person's use of the information in the Formulary.

[back to top](#)